Published online 2017 October 15.

Research Article



# Heavy Metals Contamination and Risk Assessment of Surface Soils of Babol in Northern Iran

# Abdoliman Amouei, Ali Cherati, and Dariush Naghipour<sup>3,\*</sup>

<sup>1</sup>Environmental Health Research Center (EHRC), Department of Environmental Health, Babol University of Medical Sciences, IR Iran

Received 2016 August 03; Revised 2016 December 09; Accepted 2017 January 13.

#### **Abstract**

**Background:** Currently there is much concern in the field of soil contamination with heavy metals due to rapid urbanization. Soil contamination significantly reduces environmental quality and affects human health.

**Objectives:** This study was performed to determine the level of As, Cd, Co, Cr, Ni, Pb, and Zn in the soils of Babol city in Mazandaran province and to evaluate the pollution indices of heavy metals.

**Methods:** Concentrations of heavy metals in 50 samples of surface soils (5 to 15 cm) in urban and rural areas of Babol city were measured using inductively coupled plasma optical emission spectrometry (ICP-OES). In this study, various indices including contamination factor (Cfi), degree of contamination (Cdeg), Nemerow integrated pollution index (NIPI) and the potential ecological risk (ERi) with heavy metals were assessed.

**Results:** Mean content of As, Pb, Cd, Zn, Cr, Ni, and Co in soil samples of different locations were  $6.90 \pm 2.60$ ,  $31 \pm 5.70$ ,  $0.32 \pm 0.1$ ,  $82.80 \pm 15$ ,  $32.40 \pm 8.30$ ,  $34.50 \pm 7.50$ , and  $22.6 \pm 6$  mg/kg, respectively. The average content of these metals is ranked in the following descending order: Zn > Cr > Ni > Pb > Co > As > Cd. The order of contribution of the potential ecological risk by metals was Cd (51.91%) > As (23.75) > Pb (7.61%) > Cr (5.57%) > Co (5.24%) > Zn (3.04%) > NI (2.88%).

**Conclusions:** This study indicated that the risk assessments of the soil samples with heavy metals in the studied locations are categorized from low to moderate degree of contamination.

Keywords: Heavy Metals, Surface Soil, Pollution Index, Ecological Risk Index

# 1. Background

Global industrialization and unsystematically agricultural activities have been effected natural environments and ecosystems (1, 2). Contamination of environments with heavy metals has been increased significantly after the start of the industrial revolution (3). Soil contamination with heavy metals has been reported in both developing and developed countries (4).

The metals are categorized as "heavy metals" that have a specific gravity of more than 5 g/cm³. There are well-known 60 heavy metals around the world (2). Overall, every 1000 kg of normal soil theoretically contains 200 g of chromium, 80 g nickel, 16 g lead, 0.5 g mercury, and 0.2 g cadmium (5).

Heavy metals enter human bodies by direct pathways, such as ingestion, inhalation, dermal contacts, and indirect ways or through the food chain (6). These elements may be accumulated in soils, plants, and animals, and

eventually reach human beings (7, 8). Heavy metals that are accumulated in plants could have a contradictory effect on physiological functions of plants such as photosynthesis, gaseous exchange, and nutrient absorption (6). According to previous research, heavy metals are highly hazardous to human body and could cause pathological change of organs and diseases related to the cardiovascular system, kidneys, bones, etc, and could even cause cancer owing to excessive heavy metal accumulation in human bodies (9, 10). These elements are also known to be potential inhibitors of ATPases (11). These compounds enter an organism and reach the target organ by passing through living membranes and in the process, adversely effect their permeability and degrade normal enzyme transport (12). The most dangerous heavy metals are the compounds that bind to cell membranes affecting the intracellular transport process in living forms (13).

Although heavy metals are naturally present in soil, there is metal pollution due to local sources, such as power

<sup>&</sup>lt;sup>2</sup>Agriculture and Natural Resources Center of Mazadaran Province, Mazandaran, IR Iran

<sup>&</sup>lt;sup>3</sup>Department of Environmental Health, Guilan University of Medical Sciences, IR Iran

<sup>.</sup> Corresponding author: Dariush Naghipour, Department of Environmental Health, Guilan University of Medical Sciences, IR Iran. E-mail: dnaghipour@yahoo.com

plants and iron (14, 15), steel and chemical industries (16), agriculture activities, including irrigation with polluted waters (17), sewage sludge and fertilizers (18), contaminated manure and pesticide containing heavy metals (19), waste incineration (20), combustion of fossil fuels, and road traffic (21, 22). Long-range transport of atmospheric pollutants adds to the contamination of metals in the natural environment (23).

Contamination of soils with heavy metals, especially in agricultural soils, is more important because of their potential transfer from plants to animals (feed crops) and animals to humans (food crops and vegetables) (24, 25). The use of the contaminated sludge with heavy metals, fertilizers, and animal manures could cause high contents of metals in the soils of urban and rural areas (26, 27). On the other hand, little information is available regarding human exposure and risk assessment of the soil contaminated with heavy metals in urban and rural areas in developing cities (28).

In one study by Azimzadeh et al. concentrations of heavy metals and the contamination of farmland soil samples, in 60 and 48 soil samples from urban and natural areas in center of Mazandaran province in Iran were evaluated (29). Soleimannejad et al. compared the concentrations of cadmium, zinc, copper, lead, arsenic, and iron in the soil of industrial sites and landfills of Ghaemshahr city in Mazandaran (30). Ravankhah et al. evaluated soil contamination with heavy metals in Aran and Bidgol districts of Kashan in Iran. They found that soil contamination with lead and cadmium originated from human activities, and nickel, zinc and copper were affected by natural and human factors (31).

Gong et al. studied soil pollution in rural and urban areas with potentially dangerous heavy metals (Cd, Co, Cr, Cu, Hg, Mn, Pb, Ni and Z) in the state of Wuhan, China (26). Dantu studied the situation of soil contamination in Ranga Reddy district of the southeastern part of the state of Andhra Pradesh in India to heavy metals (As, Ba, Co, Cr, Mo, Ni, Pb, Rb, Sr, V, Y, Zn, and Zr) with X-ray fluorescence spectrometry (2).

This investigation was performed to determine the level of As, Cd, Co, Cr, Ni, Pb, and Zn in collected soil samples from agricultural areas of Babol district in center of Mazandaran province and to evaluate pollution indices such as contamination factor (Cf) and degree of contamination ( $C_{\rm deg}$ ), Nemerow integrated pollution index (NIPI), and potential ecological risk (RI) in the studied locations.

#### 2. Methods

#### 2.1. Study Area

Babol city with a population of 520 000 people is the most populated city in Mazandaran province and is located in northern Iran. The city of central Mazandaran in southern Caspian Sea (15 km from Caspian Sea) is located at a longitude of 52° 44′ 20″ and latitude of 36° 24′ 15′. Average annual precipitation and temperature is 17.1°C and 799 mm, respectively. This region is one of the most important agricultural areas of Iran (29). Another characteristic of this area is the establishment of various industries and factories. The city has several rivers, especially Babolrood River that comes into the Caspian Sea.

# 2.2. Sample Collection and Preservation

Fifty soil samples in depth of 5 to 15 cm were collected from different locations of urban and rural places of Babol city (Figure 1). These composite soil samples were collected in polyethylene bags and transferred to the laboratory. The soil samples were air-dried, grounded, passed through a standard 2-mm sieve and sealed in clean bags and then stored in a refrigerator (18).

#### 2.3. Sample Preparation and Analysis

One gram of dried soil samples was digested in a digestion system (21-mL HNO3 + 7-mL HCL). The digested samples were filtered through Wathman paper (No.42) and deionized water was added to reach 50 mL and stored at 4°C. The heavy metal contents in soil samples were determined by inductively coupled plasma atomic emission spectrometry (ICP-AES), SPECTRO model, under optimum conditions. The instrument calibration was used for the quantification of selected metals and the samples (32, 33). Preparation of the reagents and calibration standards, and double distilled water was performed. The standard solutions were prepared from stock solution of 1000 mg/L by dilutions. All the soil samples were gathered with 3 replications and blank samples were analyzed simultaneously.

# 2.4. Determination of Contamination Factor (Cf) and Degree of Contamination (Cdeg)

The assessment of soil contamination is performed by Contamination factor ( $C^i_f$ ) and degree of contamination indices. The contamination factor of a metal is determined by the following equation:

$$C_f^i = \frac{C_m^i}{C_n^i} \tag{1}$$

That:

 $C_{\ m}^{i}$ , is the mean concentration of metals.




Figure 1. Soil Sampling Location Map

 $C_n^i$ , is the natural concentration of individual elements (background level).

Hakanson (34) defines four categories  $C_{\rm f}^{\rm i}$ , as follows:

a.  $C^{i}_{f} < 1$  low contamination factor => low contamination

b.  $1 < C^{i}_{f} < 3$  moderate contamination factor => mean contamination

c.  $3 < C^{i}_{f} < 6$  considerable contamination factor => high contamination

d.  $C_f^i > 6$  very high contamination factor => very high contamination

The  $C^i_f$  is a single element index, and the sum of contamination factors for all metals is indicated by the contamination degree ( $C_{deg}$ ) of the location. The degree of contamination is classified in four classes (24):

a.  $C_{deg}$  < 8 => low degree of contamination

b.  $8 < C_{deg} < 16 \Rightarrow$  moderate degree of contamination c.  $16 < C_{deg} < 32 \Rightarrow$  considerable degree of contamination

d.  $C_{\text{deg}} > 32 = very \text{ high degree of contamination}$ 

# 2.5. Evaluation of Pollution Index (PI)

The pollution level of a metal was determined by a single pollution index  $(PI_i)$ . The  $PI_i$  was calculated with the following equation:

$$PI_i = \frac{C_i}{S_i} \tag{2}$$

 $C_i$  = Metal concentration in a soil sample

 $S_i$  = Reference value

The total potential pollution of soil samples by heavy metals was evaluated by the Nemerow integrated pollution index (NIPI).

$$NIPI = \sqrt{\frac{PI_{ave}^2 + PI_{imax}^2}{2}} \tag{3}$$

Where Plave and Plimax are the average and maximum of the pollution indices for individual metals, respectively. The NIPI is categorized as follows: 1- NIPI < 0.7 => non-pollution; 2. 0.7 < NIPI <1 => warning line of pollution; 3 -1 < NIPI < 2 => low level of pollution; 4-2 < NIPI < 3 => moderate level of pollution; 5- NIPI >3 => high level pollution (26).

#### 2.6. Estimation of the Potential Ecological Risk

The potential ecological risk of heavy metal in soil samples was evaluated using the ecological risk index (RI) (27). The RI index was determined by the sum of risk factors of the metals:

$$RI = \sum_{i=1}^{7} Er^2 \tag{4}$$

Where Er<sup>i</sup> is the risk factor of any metal, and is determined as:

$$Er^{i} = Tr^{i} \times C_{f}^{i}$$

$$= Tr^{i} \frac{C_{i}}{B_{i}}$$
(5)

Ti = the toxic-response factor for any metal.

The Toxic-response values of As, Pb, Cd, Zn, Cr, Ni and Co are 10, 5, 30, 1, 2, 2 and 2, respectively.

 $C_f^i$  = the metal contamination factor calculated from the measured concentration ( $C_i$ ) and the background concentration ( $B_i$ ) of the metal.

The potential ecological risk of heavy metals in soils was classified to four categories:

a. RI < 150 => Low ecological risk

b. 150 < RI < 300 => Moderate ecological risk

c. 200 < RI < 600 => Considerable ecological risk

d. RI > 600 => Very high ecological risk

# 2.7. Statistical Analysis

Data analysis was done with the Excel software, using one-way Analysis of Variance (ANOVA) and the t-test.

# 3. Results

# 3.1. Heavy Metals Content in Soils of Selected Locations

The range of As, Pb, Cd, Zn, Cr, Ni, and Co in the soil samples were 3.1 to 12.6, 10.5 to 62.5, 0.11 to 0.5, 25.8 to 143, 20.5 to 92.6, 17.1 to 79, and 5.8 to 37.5 mg/kg, respectively (Table 1). The total concentration in the soil samples was as follows, Zn > Cr > Ni > Pb > Co > As > Cd. Overall, 80% for all samples had a content of As below 7.5 mg/kg, Pb below 36.2 mg/kg, Cd below 0.35 mg/kg, Zn below 89 mg/kg, Cr below

31 mg/kg, Ni below 35.1 mg/kg, and Co below 25.4 mg/kg. The background levels of As, Pb, Cd, Zn, Cr, Ni, and cobalt in natural and unpolluted soils of these locations were 5.2, 36.5, 0.3, 48.5, 20.8, 41.6, and 15.4, respectively. The mean contents of As, Cr, Co, and Zn in all soil samples was higher than their background levels. The mean concentrations of Cd and Pb in 40% and 20% of soil samples were higher than their background contents, respectively (Table 2).

3.2. Determination of Contamination Factor (Cf), degree of contamination (Cdeg) and Nemerow Integrated Pollution Index (NIPI) of Heavy Metals in Soil Samples of Selected Locations

Average contamination factors of selected metals in the soil samples are shown in Table 3. Among the selected metals, mean Cf of As, Pb, Cd, Zn, Cr, Ni, and Co were determined as 1.32, 0.85, 0.97, 1.70, 1.42, 0.84 and 1.34, respectively. The degree of contamination ( $C_{\rm deg}$ ) of different locations containing L1, L2, L3, L4, and L5 was 7.08, 7.82, 8.43, 9.06, and 9.57, respectively. The mean and range of Cdeg in the soil samples from different locations were 8.41 and 3.57 to 17.53, respectively.

In Table 4, the range of pollution index of Plave, Plmax, and NIPI in the soil samples of the studied locations was 0.51 to 2.50, 0.99 to 4.45 and 0.79 to 3.61, respectively. The NIPI for most of the samples (80%) showed a low level of pollution. The analysis of results showed that the average amount of the elements was in the following order, Zn (1.70) > Cr(1.42) > Co(1.34) > As(1.32) > Co(0.97) > Pb(0.85) > Ni (0.81).

3.3. Estimation of Ecological Risk of Heavy Metals in Soils of the Selected Locations

Table 5 shows the mean ecological risk factors of different metals in the soil of locations and their participation in the total potential ecological risk. Order of ecological risk of heavy metals was Cd > As > Cr > Pb > Co > NI (Figure 2). The average and range of contents of the potential ecological risk for soils of urban and rural areas was 58.48 and 22.47 to 103.31, respectively.

#### 4. Discussion

Mean content of As, Pb, Cd, Zn, Cr, Ni, and Co in soil samples of different locations were  $6.9\pm2.60$ ,  $31\pm5.70$ ,  $0.32\pm0.10$ ,  $82.80\pm15$ ,  $32.40\pm8.30$ ,  $34.50\pm7.50$ , and  $22.6\pm6$  mg.kg¹ respectively, and were larger than background levels. Total metal contents in the soil samples of different locations follow this descending order: L5 > L4 > L3 > L2 > L1. Amouei et al. determined Pb, Cd, and Zn concentrations in agricultural soils of Babol area. In this study, contents of Pb, Cd, and Zn was  $19.75\pm9.50$ ,  $0.15\pm0.1$ , and

Table 1. Heavy Metals Concentrations in the Surface Soils from the Selected Locations Location Sample As Pb Cđ Zn  $\mathbf{Cr}$ Ni Co Total <sub>HM</sub> L  $5.41 \pm 2$  $21.3 \pm 1.5$  $0.18 \pm 0.1$  $72.8 \pm 24$  $33 \pm 12.6$  $36.8 \pm 5$  $24.5 \pm 3.6$  $170 \pm 48.8$ 10  $L_2$ 10  $6.3 \pm 1.3$  $24.6 \pm 4.8$  $0.23 \pm 0.1$  $34 \pm 8.4$  $76.5 \pm 16.5$ 30.5 ± 15  $21.7 \pm 5.2$  $187.5 \pm 51$  $L_3$ 10  $7.1 \pm 2.5$  $34.4 \pm 5$  $0.30 \pm 0.1$  $83.4 \pm 12.8$  $20.5 \pm 7.4$  $209 \pm 45.4$  $32.5 \pm 10$  $35.5 \pm 7$ 10  $7.9 \pm 3.1$  $39.5 \pm 7.5$  $0.35 \pm 0.1$  $89 \pm 19.5$  $31.7 \pm 16$  $33 \pm 6.3$  $22.9 \pm 5.1$  $227.6 \pm 58$  $7.5 \pm 4.1$  $37 \pm 5.7$  $0.40 \pm 0.1$  $92.5 \pm 14$  $34 \pm 12.4$  $32.7 \pm 11$  $23.4 \pm 8.9$  $234.7 \pm 56$ Ls 10 Max 12.6 62.5 0.5 143 92.6 79 37.5 427.7 Min 3.1 10.5 0.11 25.8 20.5 17.1 82.9 22.6 Mean 6.9 31 0.32 82.8 32.4 34.5 236.6 SD 6 2.6 5.7 0.1 15 8.3 7.5 Background 36.5 48.5 41.6 168.3 5.2 0.3 20.8 15.4 P Value 0.02 0.015 0.013 0.017 0.019 0.02 0.025 0.018

Abbreviation: SD, standard deviation.

**Table 2.** Comparison of the Heavy Metals Content in the Studied Soil Samples and Maximum Allowable Contamination Based on European Union Standards

	-	
Substance (name)	Mean $\pm$ SD, mg.kg $^{ ext{-1a}}$	MAC, mg.kg <sup>-1b</sup>
Arsenic	$6.9 \pm 2.4$	5
Cadmium	$\textbf{0.32} \pm \textbf{0.1}$	1
Chrome	$32.4\pm8.3$	100
Cobalt	$22.6 \pm 6$	20
Lead	$31\pm5.7$	60
Nickel	$32.5\pm7.5$	10
Zinc	$82.8 \pm 15$	200

<sup>&</sup>lt;sup>a</sup>All data is related to the current study.

<sup>&</sup>lt;sup>b</sup>All data is belonged to European Union.




Figure 2. Percentages of Individual Metals to the Mean Potential Ecological Risk of the Soils

107  $\pm$  39.50 mg.kg-1, respectively (18). Ghorbani et al. investigated the effects of land use on the contamination of

some heavy metals (As, Pb, Cd, Cr, Cu, Fe, Ni, Se, and Zn) on the surface soils of Golestan province. The results showed that heavy metals accumulation in soil samples of industrial land was higher than agricultural and natural land (34). Dantu showed that the mean contents of As, Co, Cr, Ni, Pb, and Zn in the soils of Ranga Reddy district, Andhra Pradesh, and India were 10.5, 14.5, 67.5, 27.3, 28.3, and 77.1 mg/kg, respectively (2). Gong et al. studied the concentration of heavy metals in urban and rural top soils of Wuhan, central China. They concluded that the mean contents of Cd, Co, Cr, Pb, Ni, and Zn were 0.20, 16, 85, 33, 34, and 85 mg/kg, respectively (12). Modrzewska and Wyszkowski determined that the mean concentrations of Pb, Cd, Cr, and Ni in soils along state Road 51 leading from Olsztyn to Olsztynek, northeastern Poland, were 24.3 to 70.3, 0.31 to 0.85, 9.9 to 11.6, and 67.5 to 120 mg/kg, respectively (21). In this study, the background levels of As, Pb, Cd, Zn, Cr, Ni, and cobalt in natural and unpolluted soils of these locations were 5.2, 36.5, 0.3, 48.5, 20.8, 41.6, and 15.4, respectively. Mean contents of As, Cr, Co, and Zn in urban soil samples was larger than their background levels. The mean concentrations of Cd and Pb in 40% and 20% of soil samples were larger than their background content, respectively. Unreasonable use of pesticides and agricultural fertilizers by farmers, disposal of municipal solid wastes, and domestic and industrial wastewater sludge are the main reasons of heavy metal contamination of agricultural and urban soils in these areas (17).

Among the selected metals, mean contamination factors of As, Pb, Cd, Zn, Cr, Ni, and Co were determined as 1.32, 0.85, 0.97, 1.70, 1.42, 0.84, and 1.34, respectively. The degree of contamination ( $C_{\rm deg}$ ) of the different locations containing L1, L2, L3, L4, and L5 were 7.08, 7.82, 8.43, 9.06,

Table 3. Contamination Factor (Cf) and Degree for Heavy Metals on the Surface of Soils

Location	Sample	C <sub>fAs</sub>	CfPb	C <sub>fCd</sub>	C <sub>fZn</sub>	C <sub>fCr</sub>	C <sub>f Ni</sub>	C <sub>fCo</sub>	C <sub>deg</sub>
L <sub>1</sub>	10	1.52	0.58	0.60	1.50	1.59	0.76	0.49	7.08 <sup>a</sup>
$\mathbf{L}_{2}$	10	1.04	0.92	0.77	1.58	1.47	0.84	1.28	7.82 <sup>a</sup>
$L_3$	10	1.21	0.67	0.93	1.72	1.56	0.88	1.53	8.43 <sup>a</sup>
$L_4$	10	1.38	1.08	1.17	1.84	1.52	0.70	1.75	9.06 <sup>b</sup>
L <sub>5</sub>	10	1.44	0.99	1.33	1.91	1.63	0.81	1.65	9.57 <sup>b</sup>
Max	-	2.42	1.71	1.67	2.95	4.45	1.90	2.43	17.53
Min	•	0.60	0.29	0.37	0.53	0.99	0.41	0.38	3.57
Mean	•	1.32	0.85	0.97	1.70	1.55	0.83	1.47	8.70
SD	-	0.52	0.35	0.38	0.6	0.55	0.33	0.58	3.2

Abbreviation: SD, standard deviation.

 $\textbf{Table 4.} \ \textbf{The Pollution Indices of Heavy Metals in Soils from Selected Locations}$ 

Location	PI <sub>As</sub>	PI <sub>Pb</sub>	PI <sub>Cd</sub>	PI <sub>Zn</sub>	PI <sub>Cr</sub>	PI <sub>Ni</sub>	PI <sub>Co</sub>	PI <sub>ave</sub>	PI <sub>max</sub>	NIPI
L1	1.52	0.58	0.60	1.50	1.63	0.76	0.49	1.01	1.63	1.36 <sup>a</sup>
L2	1.04	0.92	0.77	1.58	1.39	0.84	1.28	1.12	1.58	1.37 <sup>a</sup>
L3	1.21	0.67	0.93	1.72	1.49	0.88	1.53	1.20	1.72	1.48 <sup>a</sup>
L4	1.38	1.08	1.17	1.84	1.14	0.70	1.75	1.37	1.75	1.57 <sup>a</sup>
L5	1.44	0.99	1.33	1.91	1.44	0.81	1.65	1.37	1.91	1.66 <sup>a</sup>
Max	2.42	1.71	1.67	2.95	4.45	1.90	2.43	2.50	4.45	3.61
Min	0.60	0.29	0.37	0.53	0.99	0.41	0.38	0.51	0.99	0.79
Mean	1.32	0.85	0.97	1.70	1.42	0.81	1.34	1.20	1.70	1.47
SD	0.52	0.35	0.38	0.6	0.55	0.33	0.58	3.2	0.65	0.54

Abbreviation: SD, standard deviation.

<sup>&</sup>lt;sup>a</sup>Low level of pollution.

Location	r.i	r.i	r.i	r.i	r_i	r.i	r_i		P!-I-P
Location	Er¹ <sub>As</sub>	Er¹ <sub>Pb</sub>	Er <sup>i</sup> Cd	Er¹ <sub>Zn</sub>	Er¹ <sub>Cr</sub>	Er <sup>i</sup> Ni	Er¹ <sub>Co</sub>	RI	Risk Degree
L1	15.19	2.91	18	1.50	3.18	1.53	2.27	44.67	Low
L2	10.38	4.60	23	1.58	2.94	1.69	2.56	46.60	Low
L3	12.12	3.34	1.72	28	3.12	1.75	3.05	52.96	Low
L4	13.85	5.41	35	1.84	3.04	1.36	3.49	63.23	Low
L5	14.42	4.96	40	1.91	3.26	1.39	3.30	68.86	Low
Max	24.23	8.56	50	2.95	8.90	3.8	4.87	103.31	-
Min	5.96	1.44	11	0.53	1.97	0.82	0.75	22.47	-
Mean	13.27	4.25	29	1.70	3.11	1.61	2.93	55.8	-
SD*	6.52	1.35	5.35	0.52	1.23	0.87	1.18	11.26	-

Abbreviation: SD, standard deviation.

and 9.57, respectively. In the current study, 40% of soil samples (L1, L2), based on contamination factor ( $C_f$ ) and degree of contamination ( $C_{\rm deg}$ ), were found at low levels and the others (L3, L4, L5) were categorized at a medium degree. Sayadi et al. studied the surface soils of Amir-Abad of Birjand, based on pollution and ecological indices. The results showed that the ecological risk of surface soil for the users of road-residential areas was high and the agricultural land use had moderate ecological risk and dairy farm had low ecological risk (35). Iqbal et al. showed that the average  $C_{\rm deg}$  values for the selected metal contents in urban soils from Islamabad, Pakistan during summer and winter

were 16.67 and 20.50, respectively, indicating a considerable degree of contamination (33). The NIPI for most of the samples (80%) showed a low level of pollution. The analysis results showed that the NIPI average for the elements was as follows, Zn(1.70) > Cr(1.42) > Co(1.34) > As(1.32) > Co(0.97) > Pb(0.85) > Ni(0.81). The maximum NIPI value (3.61) belonged to soil samples from urban areas and waste disposal/ treatment sites (26). The average and range of the potential ecological risk, including for soils of agricultural locations, calculated as the sum of the mean risk factors of the metals, was 58.48 and 22.47 to 103.31, respectively indicating an overall low ecological risk posed by heavy metals.

<sup>&</sup>lt;sup>a</sup>Low degree of contamination.

<sup>&</sup>lt;sup>b</sup>Moderate degree of contamination.

In the study of Jiang et al., the potential ecological risk in the order of Cd > Pb > Cu > Cr > Zn was obtained, which showed that Cd was the most important factor leading to risk (27).

#### 4.1. Conclusions

Soil contamination and the potential risk assessment with heavy metals in Babol city of Mazandaran province was investigated and 50 surface samples (5 to 15 cm) were taken from a wide range of exposed either bare soil or vegetated locations. Topsoil samples collected from urban areas (L<sub>3</sub>, L<sub>4</sub> and L<sub>5</sub>) had high As, Cd, Pb, and Zn concentrations coupled with high coefficients of variation, indicating the dominance of anthropogenic sources, where Co, Cr, and Ni exhibited generally low coefficients of variation and quite homogenous distributions across the study area, suggesting a major natural origin. Mean content of As, Pb, Cd, Zn, Cr, Ni, and Co in soil samples of different locations were  $6.9 \pm 2.60$ ,  $31 \pm 5.70$ ,  $0.32 \pm 0.10$ ,  $82.80 \pm 15$ ,  $32.40 \pm$ 8.30, 34.50  $\pm$  7.50, and 22.6  $\pm$  6 mg.kg<sup>-1</sup> respectively, and were larger than their background levels. Application of the index of contamination factor ( $C_{f}^{i}$ ), degree of contamination (C<sub>deg</sub>), Nemerow integrated potential index (NIPI), and the potential ecological risk (ER) clearly indicates that all of the soil samples were categorized in low to moderate degree of contamination and the potential ecological risk by heavy metals. The contribution order of ecological risk of heavy metals was as follows, Cd (51.91%) > As (23.75) > Pb (7.61%) > Cr(5.57%) > Co(5.24%) > Zn(3.04%) > NI(2.88%).

# Acknowledgments

The authors gratefully acknowledge food health and chemistry laboratory personnel of Guilan University of Medical Sciences for carrying out the heavy metals analysis.

#### References

- Moor C, Lymberopoulou T, Dietrich VJ. Determination of Heavy Metals in Soils, Sediments and Geological Materials by ICP-AES and ICP-MS. *Microchimica Acta*. 2001;136(3-4):123-8. doi: 10.1007/s006040170041.
- Dantu S. Heavy metals concentration in soils of southeastern part of Ranga Reddy district, Andhra Pradesh, India. *Environ Monit Assess*. 2009;149(1-4):213–22. doi: 10.1007/s10661-008-0195-8. [PubMed: 18338232].
- Kamani H, Ashrafi SD, Isazadeh S, Jaafari J, Hoseini M, Mostafapour FK, et al. Heavy metal contamination in street dusts with various land uses in Zahedan, Iran. Bull Environ Contamin Toxicol. 2015;94(3):382–6.
- Biasioli M, Barberis R, Ajmone-Marsan F. The influence of a large city on some soil properties and metals content. Sci Total Environ. 2006;356(1-3):154-64. doi: 10.1016/j.scitotenv.2005.04.033. [PubMed: 15941578].

- Vrscaj B, Poggio L, Marsan FA. A method for soil environmental quality evaluation for management and planning in urban areas. *Landscape Urban Plan*. 2008;88(2-4):81-94. doi: 10.1016/j.landurbplan.2008.08.005.
- Wang M, Markert B, Chen W, Peng C, Ouyang Z. Identification of heavy metal pollutants using multivariate analysis and effects of land uses on their accumulation in urban soils in Beijing, China. *Environ Monit Assess*. 2012;184(10):5889–97. doi: 10.1007/s10661-011-2388-9. [PubMed: 22068310].
- Onder S, Dursun S, Gezgin S, Demirbas A. Determination of Heavy Metal Pollution in Grass and Soil of City Centre Green Areas (Konya, Turkey). Polish | Environ Stud. 2007;16(1).
- Bazrafshan E, Mostafapour FK, Esmaelnejad M, Ebrahimzadeh GR, Mahvi AH. Concentration of heavy metals in surface water and sediments of Chah Nimeh water reservoir in Sistan and Baluchestan province, Iran. Desalinat Water Treat. 2016;57(20):9332–42.
- Wuana RA, Okieimen FE. Heavy metals in contaminated soils: a review of sources, chemistry, risks and best available strategies for remediation. Isrn Ecol. 2011;2011.
- Tahar K, Keltoum B. Effects of Heavy Metals Pollution in Soil and Plant in the Industrial Area, West ALGERIA. J Korean Chem Soc. 2011;55(6):1018–23. doi: 10.5012/jkcs.2011.55.6.1018.
- Bhagure GR, Mirgane SR. Heavy metal concentrations in groundwaters and soils of Thane Region of Maharashtra, India. *Environ Monit Assess*. 2011;173(1-4):643–52. doi: 10.1007/s10661-010-1412-9. [PubMed: 20300835].
- Gong M, Wu L, Bi XY, Ren LM, Wang L, Ma ZD, et al. Assessing heavy-metal contamination and sources by GIS-based approach and multi-variate analysis of urban-rural topsoils in Wuhan, central China. *Environ Geochem Health*. 2010;32(1):59–72. doi: 10.1007/s10653-009-9265-2. [PubMed: 19499347].
- Zeng X, Li LF, Mei XR. Heavy Metal Content in Chinese Vegetable Plantation Land Soils and Related Source Analysis. Agricultur Sci China. 2008;7(9):1115–26. doi: 10.1016/s1671-2927(08)60154-6.
- 14. Velea T, Gherghe L, Predica V, Krebs R. Heavy metal contamination in the vicinity of an industrial area near Bucharest. *Environ Sci Pollut Res Int.* 2009;**16 Suppl 1**:S27–32. doi: 10.1007/s11356-008-0073-5. [PubMed: 19101745].
- Shu Y, Zhai S. Study on soil heavy metals contamination of a lead refinery. Chinese J Geochem. 2014;33(4):393-7. doi: 10.1007/s11631-014-0703-1.
- Srinivasa Gowd S, Ramakrishna Reddy M, Govil PK. Assessment of heavy metal contamination in soils at Jajmau (Kanpur) and Unnao industrial areas of the Ganga Plain, Uttar Pradesh, India. J Hazard Mater. 2010;174(1-3):113–21. doi: 10.1016/j.jhazmat.2009.09.024. [PubMed: 19837511].
- Kanmani S, Gandhimathi R. Assessment of heavy metal contamination in soil due to leachate migration from an open dumping site. Appl Water Sci. 2012;3(1):193–205. doi: 10.1007/s13201-012-0072-z.
- Amouei AI, Yousefi Z, Mahvi AH, Naddafi K, Tahmasbizadeh M. Heavy Metal Concentrations in Industrial, Agricultural, and Highway Soils in Northern Iran. Environ Justice. 2012;5(3):153-7. doi: 10.1089/env.2011.0038.
- Yu S, Zhu Y, Li X. Trace metal contamination in urban soils of China. Sci Total Environ. 2012;421:17–30.
- Hu Y, Liu X, Bai J, Shih K, Zeng EY, Cheng H. Assessing heavy metal pollution in the surface soils of a region that had undergone three decades of intense industrialization and urbanization. *Environ Sci Pollut Res Int.* 2013;20(9):6150–9. doi: 10.1007/s11356-013-1668-z. [PubMed: 23546857].
- Modrzewska B, Wyszkowski M. Trace metals content in soils along the state road 51 (northeastern Poland). Environ Monit Assess. 2014;186(4):2589–97. doi: 10.1007/s10661-013-3562-z. [PubMed: 24395551].
- 22. Mico Llopis C, Peris M, Sánchez J, Recatalá L. Heavy metal content of agricultural soils in a Mediterranean semiarid area: the Segura River Valley (Alicante, Spain). *Spanish J Agric Res.* 2006;**4**(4):363. doi: 10.5424/sjar/2006044-213.

- Karanlik S, Agca N, Yalcin M. Spatial distribution of heavy metals content in soils of Amik Plain (Hatay, Turkey). Environ Monit Assess. 2011;173(1-4):181-91. doi: 10.1007/s10661-010-1380-0. [PubMed: 20221796].
- 24. Nabulo G, Young SD, Black CR. Assessing risk to human health from tropical leafy vegetables grown on contaminated urban soils. *Sci Total Environ*. 2010;**408**(22):5338–51. doi: 10.1016/j.scitotenv.2010.06.034. [PubMed: 20739044].
- dos Santos SN, Alleoni LR. Reference values for heavy metals in soils of the Brazilian agricultural frontier in Southwestern Amazonia. Environ Monit Assess. 2013;185(7):5737–48. doi: 10.1007/s10661-012-2980-7. [PubMed: 23138418].
- Qingjie G, Jun D, Yunchuan X, Qingfei W, Liqiang Y. Calculating Pollution Indices by Heavy Metals in Ecological Geochemistry Assessment and a Case Study in Parks of Beijing. *J China Univ Geosci*. 2008;19(3):230–41. doi:10.1016/s1002-0705(08)60042-4.
- Jiang X, Lu WX, Zhao HQ, Yang QC, Yang ZP. Potential ecological risk assessment and prediction of soil heavy-metal pollution around coal gangue dump. Nat Hazards Earth System Sci. 2014;14(6):1599–610. doi: 10.5194/nhess-14-1599-2014.
- Alfaro MR, Montero A, Ugarte OM, do Nascimento CW, de Aguiar Accioly AM, Biondi CM, et al. Background concentrations and reference values for heavy metals in soils of Cuba. Environ Monit Assess.

- 2015;**187**(1):4198. doi: 10.1007/s10661-014-4198-3. [PubMed: 25504190].
- Azimzadeh B, Khademi H. Estimation of background concentration of selected heavy metals for pollution assessment of surface soils of Mazandaran province, Iran. J Water Soil. 2013;27(3):548–59.
- Soleimannejad Z, Abdolzadeh A, Sadeghipour HR. Heavy Metal Concentrations in Industrial Area Soils and Landfill Site, Ghaemshahar, Iran. J Mazandaran Univ Med Sci. 2016;26(136):196-201.
- 31. Ravankhah N, Mirzaei R, Masoum S. Evaluation of Geoaccumulation Index, Contamination Factor, and Principal Component Analysis for Estimating Soil Contamination. *Iran | Health Environ*. 2015;8(3):345–56.
- AWWA; APHA. WEF (1998) Standard methods for the examination of water and wastewater. Washington: American Public Health Association;
- Iqbal J, Shah MH. Distribution, correlation and risk assessment of selected metals in urban soils from Islamabad, Pakistan. J Hazard Mater. 2011;192(2):887–98. doi: 10.1016/j.jhazmat.2011.05.105. [PubMed: 21724324].
- 34. Ghorbani H, Hafezi Moghads N, Kashi H. Effects of Land Use on the Concentrations of Some Heavy Metals in Soils of Golestan Province, Iran. J Agric Sci Technol. 2015;17(4):1025–40.
- Sayadi MH, Shabani M, Ahmadpour N. Pollution Index and Ecological Risk of Heavy Metals in the Surface Soils of Amir-Abad Area in Birjand City, Iran. Health Scope. 2015;4(1). doi: 10.17795/jhealthscope-21137.