



Reference Interval for Plasma Free Amino Acids in Iranian Children Using Reverse Phase High Performance Liquid Chromatography

Maryam Monajemzadeh,^{1,2} Sedigheh Shams,^{2,3,*} Shahrzad Mahdavi Izadi,² Seerat Elahi,² Mohammad Taghi Haghi Ashtiani,^{1,2} and Bahar Ashjaei^{2,4}

¹Department of Pathology, Tehran University of Medical Sciences, Tehran, IR Iran

²Children's Medical Center, Pediatrics Center of Excellence, Tehran, IR Iran

³Pediatric Urology Research Center, Tehran University of Medical Sciences, Tehran, IR Iran

⁴Department of Surgery, Tehran University of Medical Sciences, Tehran, IR Iran

*Corresponding author: Sedigheh Shams, Pediatrics Center of Excellence, Children's Medical Center, Dr. Qarib St, Keshavarz Blvd, Tehran, IR Iran. Tel: +98-2161472471, E-mail: shams@sina.tums.ac.ir

Received 2017 September 18; Revised 2018 January 02; Accepted 2018 February 15.

Abstract

Background: Brain damage and mental retardation in children can be caused by either excess or depletion of certain amino acids. Up to now there is no reliable reference interval for free plasma amino acid in Iranian children.

Objectives: The aim of this study was to provide the reference intervals of free plasma amino acid in healthy Iranian children using high performance liquid chromatography (HPLC) as an accurate method.

Methods: 133 children referred to Children's Medical Center, Tehran, Iran for the annual checkup entered the study after filling out the informed consent form. Demographic data and blood samples after overnight fasting were collected. Plasma amino acid concentrations were determined by reverse phase HPLC (RP-HPLC) equipped with a fluorescence detector after precolumn derivatization of amino acids with o-phthalaldehyde (OPA).

Results: Female children exhibited significantly higher plasma concentrations of taurine and lysine than male children. Other 20 amino acid values were not significantly different between two gender groups. Positive correlation was observed between age and taurine, leucine and lysine.

Conclusions: Our data provided reference intervals of free plasma amino acids in healthy Iranian children. We believe our data can be used as a guideline for pediatricians to identify the significantly deviated amino acid values in children with medical issues such as genetic disorders and some acquired illnesses. Our data is also applicable for further investigations in the field of family medicine.

Keywords: Amino Acid, High Performance Liquid Chromatography, Reference Intervals, Children

1. Background

Excess or depletion of certain amino acids will result in severe illnesses (e.g. Phenylketonuria and Maple Syrup Urine Disease) which are usually accompanied by brain damage and mental retardation (1-3). The amino acid baseline concentrations are used in assessment of nutritional status and creating amino acid level profiles (4-6). Recent studies have emphasized the importance of these nutritional profiles in prognosis of malnourished patients with chronic renal and liver diseases (7, 8). In the recent years, obtaining the nutritional profile in patients undergoing dialysis for the first time or transplant surgery, is considered as a major factor in prognosis, survival rate, mortality and morbidity (4, 8, 9).

Defining the reference interval of amino acids is important for diagnostic and prognostic purposes. Free

plasma amino acid concentrations can differ in normal individuals over a broad range of variables like diet, time of the day, gender, obesity, stress, hormones like testosterone and age. Among the identified variables, age and gender are the two main physiologic factors affecting plasma amino acids values. Significant differences between male and female and between old and young adults have been reported by several studies (4, 10-13).

Although free plasma amino acid values in children are documented in medical textbooks and journals (12, 14, 15) no equivalent study has been conducted in the Iranian child population.

2. Objectives

To determine the normal free plasma values which are the primary requirement for physicians to apply to their

practices, we have measured free plasma concentration of 22 amino acids in healthy Iranian children aged 2 to 8 years, analyzed by o-phthalaldehyde derivative high performance liquid chromatography (OPA-HPLC) method.

3. Methods

A total of 133 Iranian children (54 females and 79 males, 2-8 years old) who were referred to Children's Medical Center Hospital, Tehran, Iran for their annual check-up from July 2011 to June 2012 were recruited into the project after filling out the informed consent form by parents. The participants fulfilled the requirements of the inclusion criteria. The study size was reckoned according to CLSI guideline (C28-A2) recommendation for establishment of reference interval (16).

The inclusion criteria were as follows: a) no history of metabolic disorders, b) no history of liver or kidney diseases, c) normal development and growth d) being on a regular diet and not consuming infant formula, and finally d) no evidence of malnutrition.

Overnight fasting blood samples from antecubital vein were collected in a heparinized syringe and immediately separated by centrifugation at 3000 rpm for 15 minutes. 200 μ L of plasma was mixed with 50 μ L internal standard homoserine, afterwards ice-cold methanol (800 μ L) was applied to the mixture while vortexing, then kept in 4°C for 5 minutes.

The tubes were centrifuged at 4,000 rpm for 5 minutes and supernatant was collected. 250 μ L of supernatant was mixed with 100 μ L of borate buffer followed by 50 μ L of OPA/2ME reagent (Tube A). 200 μ L of mobile phase and 25 μ L of hydrochloric acid (HCl) was added to 50 μ L of Tube A solution and injected to HPLC system (Knauer, Germany). Separation of derivatives of amino acids was performed by reversed-phase HPLC chromatography on a C18 column using a gradient of 0.05 M acetate buffer and methanol, pH 7.02 with a flow rate of 1.0 mL per minute.

HPLC was equipped with a fluorescence detector operating at excitation wavelength of 330 nm and emission wavelength of 450 nm.

Amino acids including alanine, arginine, asparagine, aspartic acid, citrulline, α -aminobutyric acid, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, ornithine, phenylalanine, serine, taurine, threonine, tryptophane, tyrosine and valine were quantified using appropriate calibrators (Clin Cal, RECIPE, Gm pH, Germany).

Plasma amino acid concentrations were expressed as mean, 95 per cent confidence interval and scored at 5th, 25th, 75th and 95th percentiles. Amino acid concentrations were compared according to age and gender by anal-

ysis of variance (ANOVA) and t-test, respectively. IBM SPSS Statistics 17.0 software was used for statistical analysis. P value < 0.05 was considered statistically significant. The study was performed in agreement with Helsinki declaration and was conducted in accordance with considerations recommended by local ethics review committee of Tehran University of Medical Sciences (ethics code: 90-02-30-12879).

4. Results

The mean age of 133 children (54 female and 79 male) was 4.2 years (ranged from 2 to 8 years). Mean of each measured free plasma amino acid concentration, 95 per cent Confidence Interval and the scores of 5th, 25th, 75th and 95th percentiles are presented in Tables 1 and 2, respectively.

A significant correlation between age and taurine ($r = 0.185$, $P = 0.033$), leucine ($r = 0.184$, $P = 0.034$) and lysine ($r = 0.175$, $P = 0.043$) was observed.

5. Discussion

This study presents the physiologic values of 22 amino acids in Iranian children aged 2 to 8 years and their distribution as the scores of 5th, 25th, 75th and 95th percentiles. Since maximum and minimum values could have unusually high or unusually low values, we preferred to use the 95 per cent confidence interval to describe our data. These findings are intended to act as a reference for pediatricians to identify the significantly deviated amino acid values in Iranian children with medical issues such as genetic disorders and some acquired illnesses.

In the present study we used a RP-HPLC method involving pre-column derivatization with o-phthalaldehyde (OPA) to analyze the free plasma amino acid concentrations. Although simplicity and reproductivity of ion exchange chromatography (IEC) in combination with post-column ninhydrin make it the reference method for measuring the free plasma amino acids in clinical laboratories (4, 17-19), its long analytical run time made the OPA-HPLC a useful alternative method for analyzing the amino acids in physiological fluids because of OPA-HPLC's higher sensitivity and speed, both of which are very important for newborn screening (18).

The data of the present study demonstrated two additional findings. First, the plasma concentration of taurine and lysine was significantly higher in female children than in male ones and the other 20 measured amino acids were not significantly different between the two gender groups. Our results were in consistency with the report of Gregory

Table 1. Mean Concentration and 95% Confidence Interval of Amino Acids in Healthy Children

Amino Acid	Mean, $\mu\text{Mol/L}$	Confidence Interval 95%	Amino Acid	Mean, $\mu\text{Mol/L}$	Confidence Interval 95%
Alanine	376.39	358.36 - 402.59	Leucine	115.379	108.48 - 122.24
Arginine	71.813	67.10 - 76.15	Lysine	131.099	123.46 - 136.07
Asparagine	55.4815	53.25 - 57.99	Methionine	26.44	24.49 - 28.75
Aspartic acid	5.92	5.34 - 6.35	Ornithine	62.02	58.42 - 64.83
Citrulline	26.52	24.97 - 28.04	Phenylalanine	61.40	59.00 - 64.49
γ -Aminobutyric acid	23.10	21.53 - 24.25	Serine	139.68	134.20 - 144.07
Glutamic acid	59.71	54.21 - 65.27	Taurine	77.11	68.11 - 82.61
Glutamine	552.12	534.19 - 566.89	Threonine	121.18	113.61 - 127.00
Glycine	258.15	245.97 - 270.03	Tryptophane	50.15	47.20 - 53.22
Histidine	91.56	86.71 - 95.14	Tyrosine	67.01	63.26 - 70.57
Isoleucine	66.72	62.72 - 70.90	Valine	227.48	216.76 - 238.17

Table 2. Values of Amino Acids for the Scores of 5th, 25th, 75th and 95th Percentiles

Amino Acid	Percentile, $\mu\text{Mol/L}$			
	5	25	75	95
Alanine	184.06	284.50	466.20	581.16
Arginine	37.02	55.70	82.40	134.08
Asparagine	35.82	45.10	64.70	79.52
Aspartic acid	2.36	3.70	7.30	11.58
Citrulline	13.70	20.20	32.60	42.10
γ -aminobutyric acid	13.22	17.60	25.40	41.78
Glutamic acid	22.44	35.87	76.77	127.95
Glutamine	372.88	495.00	628.80	687.00
Glycine	164.62	208.80	308.40	408.40
Histidine	56.340	73.90	108.40	135.44
Isoleucine	37.020	50.10	77.00	114.42
Leucine	65.22	88.90	134.30	196.50
Lysine	85.26	102.10	148.80	202.34
Methionine	11.72	18.00	31.70	50.16
Ornithine	33.02	48.70	73.40	94.88
Phenylalanine	38.91	50.75	71.13	97.87
Serine	95.34	118.00	154.60	194.00
Taurine	27.64	44.40	98.80	178.72
Threonine	64.94	93.10	140.20	197.00
Tryptophane	15.62	40.10	61.00	79.70
Tyrosine	38.72	51.90	82.00	110.52
Valine	144.40	181.20	264.10	346.86

et al that the effect of sex was not apparent in children, although they found significant difference for 5 amino acids

including isoleucine, leucine, methionine, phenylalanine, and tyrosine in adolescents (20). Their values were significantly lower in adult females than in males. In a study conducted by Yamamoto et al. a link between sex and concentration of most of the amino acids especially branched chain amino acids was reported (21). A few other studies have also identified lower concentrations of branched-chain amino acids in older females (10, 11). Since several studies found sex-dependent amino acid values in adolescents and no significant differences in preadolescent children (10, 22, 23), it is believed that these gender-related differences occur after adolescence.

Recent studies on amino acid concentrations have described a relation between diet and amino acid values. Milson et al studied on fasting and postprandial plasma amino acid concentration and showed a gender-related difference in fasting subjects (24). They have reported that fasting female subjects had lower values in several amino acids than fasting male ones, but this effect was insignificant in postprandial subjects. Therefore, dietary protein can induce a significant degree of variation in amino acid values throughout a day. Another study, conducted by Rana et al. showed that protein intake is significantly lower in vegan diets particularly taurine which was absent in vegan diets (25). In comparison to these data, we suggest that the lower amount of taurine and lysine could be a result of a daily diet variation in children. It needs a study with more cases and precise diet control.

Our second finding identified a significant relation between age and lysine, leucine and taurine values. Our results to some extent, were consistent with previous studies which are suggesting an increase in amino acid levels in growing children probably caused by increasing muscle mass (26). Lepage et al. demonstrated a unique pattern of age-specific distribution for each amino acid in a healthy pediatric population (27). In their study, alanine, arginine, asparagine, methionine, ornithine, phenylalanine, proline, threonine and tyrosine were decreased in the first years of life, and then increased steadily up to 18 years of age. An Initial reduction followed by stable concentration was observed in taurine and serine. Aspartic acid and glutamic acid had decreasing trend in all ages. Citrulline demonstrated a two-step increment at 0 to 3 years of age and 13 to 15 years. Cystine, glutamine, glycine, histidine, isoleucine, leucine, lysine, tryptophan, and valine showed increasing values throughout 0 to 18 years of age. Several other studies (20, 28, 29) have also reported a specific pattern of age-dependent amino acid concentrations with slight differences to Lepage et al.'s study. It seems that children under one year need specific reference value. Cruz et al. in their study validated the HPLC method for determination plasma amino acids and established refer-

ence interval for Brazilian population (30). They recommended that each laboratory should establish its own reference interval due to variation in methodology and analytical techniques used in published data. The main limitation of our study, as that of Cruz et al.'s, was absence of specimens from children under 2 years old. That was because their parents refused to give permission for blood collection from their babies for our study.

This study provided reference intervals of free plasma amino acids in healthy children. We believe our data can be used as a guideline for investigations in the field of family medicine and in the practice of pediatricians. We have also reported an significant difference between age and 3 of amino acid levels and also a significant higher concentration of taurine and lysine in female children whereas other amino acids were not significantly different, but it is more ideal to establish further follow up studies and include more affecting factors such as, body mass index, and diet.

Acknowledgments

This research has been supported by Tehran University of Medical Sciences. We thank all staff of clinical laboratory of children's medical center specially Mr. Mohsen Sivashi and Mrs. Neda Rezaei for specimen collection, processing and analyzing.

Footnote

Funding/Support: This research project was funded by Tehran University of Medical Sciences.

References

1. Svasti J, Wasant P, Tiensuwan M, Sawangareetrakul P, Srisomsap C, Pangkanon S, et al. Normal plasma free amino acid levels in Thai children. *J Med Assoc Thai*. 2001;**84**(11):1558-68. [PubMed: [11853298](#)].
2. Lin G, Wang X, Wu G, Feng C, Zhou H, Li D, et al. Improving amino acid nutrition to prevent intrauterine growth restriction in mammals. *Amino Acids*. 2014;**46**(7):1605-23. doi: [10.1007/s00726-014-1725-z](#). [PubMed: [24658999](#)].
3. Burrage LC, Nagamani SC, Campeau PM, Lee BH. Branched-chain amino acid metabolism: from rare Mendelian diseases to more common disorders. *Hum Mol Genet*. 2014;**23**(R1):R1-8. doi: [10.1093/hmg/ddu123](#). [PubMed: [24651065](#)].
4. Corte Z, Venta R. Biological variation of free plasma amino acids in healthy individuals. *Clin Chem Lab Med*. 2010;**48**(1):99-104. doi: [10.1515/CCLM.2010.008](#). [PubMed: [19929754](#)].
5. Rudman D, Mattson DE, Feller AG, Cotter R, Johnson RC. Fasting plasma amino acids in elderly men. *Am J Clin Nutr*. 1989;**49**(3):559-66. [PubMed: [2923089](#)].
6. Palova S, Charvat J, Masopust J, Klapkova E, Kvapil M. Changes in the plasma amino acid profile in anorexia nervosa. *J Int Med Res*. 2007;**35**(3):389-94. doi: [10.1177/147323000703500314](#). [PubMed: [17593868](#)].

7. Ivarsen P, Tietze IN, Pedersen EB. Nutritional status and amino acids in granulocytes and plasma in patients with chronic renal disease and varying residual renal function. *Nephron*. 2001;**88**(3):224–32. doi: [10.1159/000045994](https://doi.org/10.1159/000045994). [PubMed: [11423753](https://pubmed.ncbi.nlm.nih.gov/11423753/)].
8. Henkel AS, Buchman AL. Nutritional support in patients with chronic liver disease. *Nat Clin Pract Gastroenterol Hepatol*. 2006;**3**(4):202–9. doi: [10.1038/ncpgasthep0443](https://doi.org/10.1038/ncpgasthep0443). [PubMed: [16582962](https://pubmed.ncbi.nlm.nih.gov/16582962/)].
9. Schulte-Frohlinde E, Wagenpfeil S, Willis J, Lersch C, Eckel F, Schmid R, et al. Role of meal carbohydrate content for the imbalance of plasma amino acids in patients with liver cirrhosis. *J Gastroenterol Hepatol*. 2007;**22**(8):1241–8. doi: [10.1111/j.1440-1746.2006.04620.x](https://doi.org/10.1111/j.1440-1746.2006.04620.x). [PubMed: [17688664](https://pubmed.ncbi.nlm.nih.gov/17688664/)].
10. Caballero B, Gleason RE, Wurtman RJ. Plasma amino acid concentrations in healthy elderly men and women. *Am J Clin Nutr*. 1991;**53**(5):1249–52. [PubMed: [2021131](https://pubmed.ncbi.nlm.nih.gov/2021131/)].
11. Nakamura H, Nishikata N, Kawai N, Imaizumi A, Miyano H, Mori M, et al. Plasma amino acid profiles in healthy East Asian subpopulations living in Japan. *Am J Hum Biol*. 2016;**28**(2):236–9. doi: [10.1002/ajhb.22787](https://doi.org/10.1002/ajhb.22787). [PubMed: [26407660](https://pubmed.ncbi.nlm.nih.gov/26407660/)].
12. Armstrong MD, Stave U. A study of plasma free amino acid levels. II. Normal values for children and adults. *Metabolism*. 1973;**22**(4):561–9. [PubMed: [4696900](https://pubmed.ncbi.nlm.nih.gov/4696900/)].
13. Galante A, Angelico F, Crocchioni G, Pennetti V. Intersexual differences in the serum-free amino acid pattern of young adults, normal obese aged subjects. *Nutr Metab*. 1978;**22**(2):119–26. [PubMed: [619318](https://pubmed.ncbi.nlm.nih.gov/619318/)].
14. Babu SV, Shareef MM, Shetty AP, Shetty KT. HPLC method for amino acids profile in biological fluids and inborn metabolic disorders of aminoacidopathies. *Indian J Clin Biochem*. 2002;**17**(2):7–26. doi: [10.1007/BF02867967](https://doi.org/10.1007/BF02867967). [PubMed: [23105346](https://pubmed.ncbi.nlm.nih.gov/23105346/)].
15. Wu PY, Edwards N, Storm MC. Plasma amino acid pattern in normal term breast-fed infants. *J Pediatr*. 1986;**109**(2):347–9. [PubMed: [3734973](https://pubmed.ncbi.nlm.nih.gov/3734973/)].
16. *How to Define and Determine Reference Intervals in the Clinical Laboratory; Approved Guideline*. Wayne; 2000. Clinical Laboratory and Standards Institute. p. 13–22.
17. Le Boucher J, Charret C, Coudray-Lucas C, Giboudeau J, Cynober L. Amino acid determination in biological fluids by automated ion-exchange chromatography: performance of Hitachi L-8500A. *Clin Chem*. 1997;**43**(8 Pt 1):1421–8. [PubMed: [9267323](https://pubmed.ncbi.nlm.nih.gov/9267323/)].
18. Fekkes D. State-of-the-art of high-performance liquid chromatographic analysis of amino acids in physiological samples. *J Chromatogr B Biomed Appl*. 1996;**682**(1):3–22. [PubMed: [8832421](https://pubmed.ncbi.nlm.nih.gov/8832421/)].
19. Fekkes D, Voskuilen-Kooyman A, Jankie R, Huijman J. Precise analysis of primary amino acids in urine by an automated high-performance liquid chromatography method: comparison with ion-exchange chromatography. *J Chromatogr B Biomed Sci Appl*. 2000;**744**(1):183–8. [PubMed: [10985580](https://pubmed.ncbi.nlm.nih.gov/10985580/)].
20. Gregory DM, Sovetts D, Clow CL, Scriver CR. Plasma free amino acid values in normal children and adolescents. *Metabolism*. 1986;**35**(10):967–9. [PubMed: [3762400](https://pubmed.ncbi.nlm.nih.gov/3762400/)].
21. Yamamoto H, Kondo K, Tanaka T, Muramatsu T, Yoshida H, Imaizumi A, et al. Reference intervals for plasma-free amino acid in a Japanese population. *Ann Clin Biochem*. 2016;**53**(Pt 3):357–64. doi: [10.1177/0004563215583360](https://doi.org/10.1177/0004563215583360). [PubMed: [25829462](https://pubmed.ncbi.nlm.nih.gov/25829462/)].
22. Fernstrom JD, Wurtman RJ, Hammarstrom-Wiklund B, Rand WM, Munro HN, Davidson CS. Diurnal variations in plasma concentrations of tryptophan, tyrosine, and other neutral amino acids: effect of dietary protein intake. *Am J Clin Nutr*. 1979;**32**(9):1912–22. [PubMed: [573061](https://pubmed.ncbi.nlm.nih.gov/573061/)].
23. Lavie L, Lavie P. Daily rhythms in plasma levels of homocysteine. *J Circadian Rhythms*. 2004;**2**(1):5. doi: [10.1186/1740-3391-2-5](https://doi.org/10.1186/1740-3391-2-5). [PubMed: [15347422](https://pubmed.ncbi.nlm.nih.gov/15347422/)].
24. Milsom JP, Morgan MY, Sherlock S. Factors affecting plasma amino acid concentrations in control subjects. *Metabolism*. 1979;**28**(4):313–9. [PubMed: [571951](https://pubmed.ncbi.nlm.nih.gov/571951/)].
25. Rana SK, Sanders TA. Taurine concentrations in the diet, plasma, urine and breast milk of vegans compared with omnivores. *Br J Nutr*. 1986;**56**(1):17–27. [PubMed: [3676193](https://pubmed.ncbi.nlm.nih.gov/3676193/)].
26. Armstrong MD, Stave U. A study of plasma free amino acid levels. III. Variations during growth and aging. *Metabolism*. 1973;**22**(4):571–8. doi: [10.1016/0026-0495\(73\)90070-x](https://doi.org/10.1016/0026-0495(73)90070-x).
27. Lepage N, McDonald N, Dallaire L, Lambert M. Age-specific distribution of plasma amino acid concentrations in a healthy pediatric population. *Clin Chem*. 1997;**43**(12):2397–402. [PubMed: [9439460](https://pubmed.ncbi.nlm.nih.gov/9439460/)].
28. Janas LM, Picciano MF, Hatch TF. Indices of protein metabolism in term infants fed human milk, whey-predominant formula, or cow's milk formula. *Pediatrics*. 1985;**75**(4):775–84. [PubMed: [3872443](https://pubmed.ncbi.nlm.nih.gov/3872443/)].
29. Scott PH, Sandham S, Balmer SE, Wharton BA. Diet-related reference values for plasma amino acids in newborns measured by reversed-phase HPLC. *Clin Chem*. 1990;**36**(11):1922–7. [PubMed: [2242570](https://pubmed.ncbi.nlm.nih.gov/2242570/)].
30. Cruz AF, Barbosa TMCC, Adelino TER, Lima WP, Mendes MO, Valadares ER. Amino acid reference intervals by high performance liquid chromatography in plasma sample of Brazilian children. *J Brasileiro de Patologia e Medicina Laborat*. 2016. doi: [10.5935/1676-2444.20160012](https://doi.org/10.5935/1676-2444.20160012).